

GROWTH AND DEVELOPMENT OF *ECHINACEA PURPUREA (L). MOENCH* PLANT IN SIRDARYA REGION.**Ergasheva Nilufar Abdullaevna**

Lecturer, Department of «Medicinal plants and botany» Gulistan State University, Uzbekistan. nilufar.ergasheva16@gmail.com

Annotation: *The order of sowing the seeds of perennial herb *Echinacea purpurea L.* belonging to the family *Asteraceae* in open fields in Mirzachul region of Uzbekistan, seed germination, plant growth and information on developmental biology, flowering biology and seed yield, plant irrigation and cultivation, yield harvesting rules.*

Keywords: *Mirzachul oasis, medicinal plant, seed germination, growth and development biology, flowering biology, seed yield, biomorphological features, ontogeny, phenospector.*

Echinacea purpurea L. is native to North America, first entering European countries and then Central Asia in various ways. In some countries it has been introduced as an ornamental plant, and in some cases echinacea seeds have become widespread as a weed among the seeds of agricultural crops [1]. Today *Echinacea purpurea L.* is widely used not only in folk medicine but also in scientific medicine, its underground and surface preparations are included in the state pharmacopoeias of many European countries, the Russian Federation, Belarus and Ukraine, rich in biologically active substances is a medicinal plant [2]. *Echinacea purpurea L.* is recommended for use in patients with depression and during periods of mental and physical stress, as well as in improving the body's immune system, metabolism, colds, diabetes and liver disease, as well as disease prevention. tadi [3,4]. The surface parts of *Echinacea purpurea L.* contain polysaccharides, essential oils, flavanoids, additives, saponins, echinocside glycosides, organic acids, the roots and rhizomes of the underground parts contain inulin, glucose, essential oil, phenolic carbonic acid betaine and resins [5,6]. Given that *Echinacea purpurea L.* is a valuable medicinal plant, in order to meet the demand for raw materials in the pharmaceutical industry of Uzbekistan, it is necessary to plant this plant in different regions of the country, to study its eco-biological properties warming up [7].

Echinacea purpurea L. planted in experimental plots around Gulistan, Syrdarya region. Before sowing, the seeds were stratified overnight in an aqueous solution of ash. *Echinacea* seeds are adapted to the weather and soil conditions for planting. The seeds were sown in 2 variants. In Option 1, seeds were sown in rows on prepared plots, and in Option 2, seeds were sown in nests. 3-4 seeds were sown in each nest. It is known that in the treatment of red echinacea used above-ground and underground parts. Therefore, if option 1 provides for a large mass of the plant from the above-ground parts - stems,

leaves and flowers, then option 2, which is planted in the nest, should get more mass from the underground parts, ie rhizomes and roots designed. The distance between each nest is 10-15 cm, which allows the plant to grow better in the underground parts. The biological feature of red echinacea is that during the 1st year of vegetation, only the pre-root buds form, ending their vegetation, and the generative phase begins in the 2nd year of vegetation. At the end of the first year of vegetation, 15-20 root balls are formed, and the plant completes its vegetation and enters the winter. In the first year of vegetation, 2 seeds germinate in seeds sown in early spring and last for 35-40 days. The leaves of the plant germinate 9-10 days after sowing and perform assimilation functions together with the seeds, the average length of the seed pods is 5,5 mm and the width is 3,5 mm, the first chin the leaf is on average 10,5 mm long and 6 mm wide, with ovoid, oval edges with straight teeth. Our observations on the experimental site showed that every 10 days, the first, second and third order leaves appeared on the grass. The growth and development of plants has changed dramatically since the introduction of agro-technical measures, such as mowing, grazing, watering, fertilizing. In the first year of vegetation, weeding, mowing, watering after each mowing, feeding with local and mineral fertilizers were carried out 2-3 times a month. More local fertilizers were used in the feeding, as the use of local fertilizers (manure) for medicinal plants is of great ecological and pharmacological importance. As mentioned above, at the end of the first year of vegetation, *Echinacea purpurea* L. produces 15-20 pre-rooted leaves, in which case the plant completes its vegetation and enters the winter. During the second year of vegetation of *Echinacea purpurea* L., in both variants, the period of flower stem formation, weeding and flowering period and duration is from May 5 to June 10, flowering lasts until August 12, seed ripening It runs from August 13 to October 15 it should be noted that when *Echinacea purpurea* L. is planted in rows and nests, the height of the seedlings and the size of the flower are slightly different, so the height and flowers of echinacea in the row planting variant are slightly smaller than those planted in the nest. The main reason for this is that when planted in rows, the seeds and sprouted seedlings are placed close to each other, resulting in the seedlings are slightly smaller in height and flowers than in the nested options, the seedlings in the nested area are tall and the flowers are large, which is related to nutrient content As a result, there are more opportunities for nutrition and development in nested areas. *Echinacea* seedlings, which germinate in early April, form flowering stems in late April and early May and enter the flowering phase from May 5. The total flowering occurs in the tenth of June. The flowering period of echinacea in both variants begins in early May. corresponds to. In the experimental plot, the general flowering occurs on June 30, and in early August the flowering ends, the duration of flowering is 60-65 days. The ripening of echinacea seeds lasts from the first ten days of August to mid-October. The ripening of the seeds depends on the time of flowering of the echinacea, usually large balls of flowers appear at the main part of the stem, i.e. at the tip, and then the stalks on the lateral

branches of the stem bloom, the branches on the stems are arranged in series, the seed They ripen earlier in these first- and second-stage inflorescences and produce quality seeds. Therefore, when preparing raw materials from the surface of the echinacea, it is recommended to first collect the seeds formed in the inflorescence. Then they start collecting raw materials for medicines. It is usually advisable to harvest the raw material during the general flowering period, as this is the period when the plant has the maximum amount of biologically active substances. In order to prepare enough seeds for the next year, the farm needs to set aside separate areas. At the end of the second year of the echinacea vegetation, medicinal raw materials can be prepared from the surface of the plant, so that the seeds ripen when the plants are in full bloom or because of the farm's need for red echinacea seeds. raw materials can be prepared. In the planted area of the experimental plot, one third of each variant was allocated a test area of 1 m², the plant parts were harvested and dried separately in the shade, with an average of 3.1 kg of wet raw material from the row variant. was obtained, and the dried mass was 785 grams. The average wet yield from the nesting area was 2.9 kg per 1 m², and when dried it was 735 grams. The differences between the two types of raw materials are negligible. Calculations show that 3 tons of wet raw material can be prepared from 1 hectare of red echinacea area, which is 760 kg dry. Depending on the type of drug produced in the pharmaceutical industry, red echinacea can be used both wet and dry. Seed productivity. In order to determine the seed yield of the Red echinacea plant in the experimental field, the seeds in the plant baskets of each variant were harvested with full ripening and the average seed yield was determined. According to the data obtained, the seed yield of the medicinal echinacea plant grown in the experimental field varied in different amounts depending on the method of sowing and years of cultivation. Of the variants studied, relatively high seed yields were obtained in 3-year-old plants. The seed yield from this variant averaged 2.34 s / ha. The lowest seed yield in the experimental field was observed in 1 variant, ie in the first year when sowing from seed (0.32 s / ha). In the second variant of the experiment, 0.84 s / ha was obtained from plants sown in the first year, and 2.05 s / ha from the variant sown in the second year (3 variants). In general, by studying the seed yield of *Echinacea purpurea* L., it can be concluded that it is possible to obtain a seed yield of 2.34 s / ha higher than that of 3-year-old plants after sowing. At the first observation of seeds (July 28) the seeds are small, incomplete, light green (0.6-0.6 mm), the seeds are firmly attached to the branches, the seeds are brittle when squeezed by hand, the humidity is high It was noted that When the degree of ripening of seeds is determined on August 10 (15 days), their average size is 1-1.2 mm, brown, the seeds are lightly clinging to the branches, not crushed when squeezed with a hard hand, dry moisture level is not noticeable . When the ripening rate of the seeds was determined on August 15, the color of the seeds turned dark brown and became very hard, and most of the seeds were spilled on the branches. In general, observations on the maturation of the seeds of the Red echinacea plant showed that its seeds are fully

mature within 14-16 days after emergence. During this period, the seeds turn brown, the seeds are 1.0-1.2 mm in diameter, lightly clinging to the branches, and are not crushed when compressed, allowing the seeds to be harvested. After this period, it is observed that 30-40% of them can be spilled. In the study of seed germination of the Red echinacea plant in the experimental field, the number of flowers in the baskets during the period when the plant entered the full flowering phase, the number of seeds in the period of full seed germination and the number of seeds formed during flowering ratio (in%), weight of seeds in one plant and weight of 1000 seeds were determined. During the full flowering phase of the plant, the number of flowers in a basket increased from 80 to 258. The weight of 1000 seeds in the first year of flowering is relatively low mass - 0.75 g, in the second year of flowering the weight of 1000 seeds is relatively heavy - 0.86 g. formed. To determine the purity of the seeds of the red echinacea plant, add 4.0 g of each variant. 3 samples were taken and each seed sample was examined using a separate magnifying glass and fully ripened, all the seeds into one piece, damaged, brushed, crooked, into the second piece, the straw between the seeds, The broken seed pieces were divided into a third piece. Each seed sample was weighed on a separate scale to determine its weight and the percentage of the total sample weight (4.0 g). From the results it can be concluded that the production of abundant and high-quality seeds depends on the type of crop, the biological characteristics of the cultivated variety, the applied agro-technological measures (tillage, sowing method, duration, norms, row spacing) , fertilization, irrigation, weed, pest and disease control), timing of harvest, techniques and technologies, seed cleaning, storage. The choice of area is important in the cultivation of red echinacea.

Conclusion. Seeds of the plant *Echinacea purpurea* L. have high germination rates in laboratory and experimental conditions, with 88-95% germination recorded in all variants. It grew and developed as a spring-autumn plant in the Mirzachel oasis.

LITERATURE

1. Gomilevskiy V. I. Medicinal plants, useful for culture on the Caspian coast // *Selskiy khozyain*. – M.: 1915. – № 45. – S. 1727-1728.
2. Kshnikatkina A.N., Gushina V.A. Productivity of exinatseipurpurnoy (*Echinacea purpurea*) in dependence on regulators of growth // *Novie in traditsionnyye rasteniyaperspektivi ix ispolzovaniya: mezhdunarodniy simposium*. - M.: Izd-vo RUDN, 2001 - T. 2. - P. 314-316.
3. N. Ergasheva, Tagayeva D, Abdurashidova M. Flowering and fertilization biology of *Echinacea purpurea* L. in Mirzachel oasis // *International journal for innovative engineering and management Research*. Vol 10 Issue, May 2021.
4. Samorodov V.N., Pospelov S.V., Moiseeva G.F., Sereda A.V. Phytotoxic composition of representatives of the genus *Echinacea* (*Echinacea* Moench) and ego

pharmacological properties (review) // Xim. farm. magazine. - 1996. - T. 30, № 4. - P. 32-37

5. Zakirov P. K., Tashmukhamedov R. I., Kabulov A. R. Some perspective medicinal plants of southern Uzbekistan // VII delegates sezd VBO: Tez. dokl. –L., 1983. –C. 192-194.

6. Titova O. A. To increase the productivity of some species of the genus *Asparagus* (*Asparagus* L.), introduced in the Botanical Garden of the USSR (Introduction and acclimatization of plants: Sb. nauch. tr. –Tashkent, BS AN UzSSR, 1988. –vyp. 22. –S. 43-48.

7. Khazanovich R.A., Russian M.I., Gomolitsskiy P.A. Experience of some medicinal plants in Tashkent / Trudi Botanicheskogosada AN UzSSR: –Tashkent, BS AN UzSSR, 1951. –Vip.2. – S.181-190.